

Asymptotic expansions of unstable (stable) manifolds in time-discrete systems

Shin-itiro Goto and Kazuhiro Nozaki

Department of Physics, Nagoya University, Nagoya 464-8602, Japan

Abstract

By means of an updated renormalization method, we construct asymptotic expansions for unstable manifolds of hyperbolic fixed points in the double-well map and the dissipative Hénon map, both of which exhibit the strong homoclinic chaos. In terms of the asymptotic expansion, a simple formulation is presented to give the first homoclinic point in the double-well map. Even a truncated expansion of the unstable manifold is shown to reproduce the well-known many-leaved (fractal) structure of the strange attractor in the Hénon map.

1 Introduction

The bifurcation of separatrices and homoclinic or heteroclinic structures are well known to lead to genesis of chaos in conservative dynamical systems[1]. While numerical iterations of low-dimensional mappings easily provide these complicated structures, it is extremely difficult to derive them analytically. For nearly integrable systems, the difficulty may have been overcome by means of the asymptotic expansion beyond all orders supplemented with the Borel sum and the stokes phenomenon [2][3] [4] [5]. However, this method can not be applied to the systems far from integrable such as a finite-time-discrete dynamical system with double-well potential (the double-well map) and the strange attractor in the dissipative Hénon map, which will be analyzed in the present paper.

Recently a novel method based on the perturbative renormalization group theory has been developed as an asymptotic singular perturbation technique [6]. The renormalization group (RG) method removes secular or divergent terms

from a perturbation series by renormalizing integral constants of lower order solutions. This RG method was reformulated on the basis of a naive renormalization transformation and the Lie group [8]. The application of the RG method to some non-chaotic discrete systems was achieved in the framework of the envelope method [7]. In this paper, we apply the reformulated RG method to chaotic discrete systems. Our approach is straightforward and much simpler than the original RG method [6] or the envelope method [7] and we obtain asymptotic expansions of unstable and stable manifolds constituting homoclinic tangles in the double-well map and an unstable manifold included in the closure of the strange attractor of the dissipative Hénon map.

2 Double well map

Let us analyze a symplectic map obtained by time-discretization of canonical equations for dynamical systems with a single freedom:

$$x_{n+1} - x_n = p_{n+1}, \quad p_{n+1} - p_n = \delta^2 f(x_n). \quad (1)$$

For concreteness, we choose a double well potential

$$f(x_n) = x_n - 2x_n^3.$$

Then we get the second-order difference equation for x_n

$$Lx_n \equiv x_{n+1} - (2 + \delta^2)x_n + x_{n-1} = -2\delta^2 x_n^3, \quad (2)$$

where δ^2 represents the time difference and is not a small parameter in general. This map is area-preserving and has a hyperbolic fixed point at $(x, p) = (0, 0)$. Although we concentrate on the system with a double-well potential in this section, the following analysis will hold for more general systems such as the standard map. For $\delta^2 \rightarrow 0$ (i.e. in the continuous limit), the system becomes integrable and the phase space is occupied by regular orbits separated by a separatrix. If $\delta^2 \neq 0$, the splitting of separatrix occurs and unstable and stable manifolds of the hyperbolic fixed point cross each other at an infinite number of points (homoclinic points) that accumulate to the hyperbolic fixed point. This complex homoclinic structure (homoclinic tangles) is the typical chaos generated by the Birkhoff-Smale's horse shoe mechanism. When δ^2 is small enough, the splitting of separatrix is very small and an asymptotic expansion of unstable and stable manifolds

was obtained by means of the singular perturbation method of beyond all order [3]. Here, we are interesting in the case that $\delta^2 \sim \mathcal{O}(1)$ or larger. Let us construct a formal series solution x_n^u near the hyperbolic fixed point $(x, p) = (0, 0)$ along the unstable manifold, that is, $\lim_{n \rightarrow -\infty} x_n^u = 0$: (For brevity, x_n^u will be noted as x_n in the following.)

$$x_n = x_n^{(1)} + x_n^{(2)} + x_n^{(3)} + x_n^{(4)} + \dots \quad (3)$$

Here δ^2 is not necessarily small and $|x_n^{(1)}| > |x_n^{(2)}| > \dots$. Then, naive perturbed equations for $x_n^{(j)}$ are written as

$$\begin{aligned} Lx_n^{(1)} &= 0, & Lx_n^{(2)} &= 0, \\ Lx_n^{(3)} &= -2\delta^2 x_n^{(1)3}, & Lx_n^{(4)} &= -6\delta^2 x_n^{(1)2} x_n^{(2)}, \\ &\dots \end{aligned}$$

The leading-order solution satisfying the condition $\lim_{n \rightarrow -\infty} x_n = 0$ is given by

$$x_n^{(1)} = AK^n, \quad (4)$$

where A is an integral constant, $\delta = 2 \sinh(k/2) > 0$ and $K = \exp k > 1$. The series solution (3) is obtained as

$$x_n = AK^n \left(1 - c_2 A^2 K^{2n} + c_4 A^4 K^{4n} - c_6 A^6 K^{6n} + \dots \right), \quad (5)$$

where c_2 , c_4 and c_6 are constants defined as

$$\begin{aligned} c_2 &= \frac{\delta^2}{D_3}, & c_4 &= \frac{3\delta^4}{D_3 D_5}, \\ c_6 &= \frac{3\delta^6}{D_7 D_3} \left(\frac{1}{D_3} + \frac{3}{D_5} \right), & D_j &= \cosh jk - \cosh k \quad (j \in \mathbf{N}). \end{aligned}$$

Now we introduce a renormalized variable $\tilde{A}(n)$ so that all the higher order terms in the series solution (5) are renormalized into the integral constant A , that is,

$$x_n = \tilde{A}(n) K^n, \quad (6)$$

where

$$\tilde{A}(n) = A \left(1 - c_2 A^2 K^{2n} + c_4 A^4 K^{4n} - c_6 A^6 K^{6n} + \dots \right), \quad (7)$$

which is called a renormalizaion transformation $A \rightarrow \tilde{A}$ [8]. In order to derive a difference equation (a renormalizaion equation) for the renormalized

variable $\tilde{A}(n)$, we express a difference $\tilde{A}(n+1) - \tilde{A}(n)$ in terms of $\tilde{A}$. From (7), we have

$$\begin{aligned}\tilde{A}(n+1) - \tilde{A}(n) &= -c_2(K^2 - 1)K^{2n}A(n)^3 \\ &\quad + c_4(K^4 - 1)K^{4n}A(n)^5 + \dots\end{aligned}\quad (8)$$

By solving (7) with respect to A iteratively, A can be expressed in terms of $\tilde{A}$ and n as

$$A = \tilde{A}(n)\{1 + c_2\tilde{A}(n)^2K^{2n} + (3c_2^2 - c_4)\tilde{A}(n)^4K^{4n} + \dots\}.$$

Thus we can replace A in (8) by $\tilde{A}$ and obtain a renormalizaion equation:

$$\begin{aligned}\tilde{A}(n+1) - \tilde{A}(n) &= -c_2(K^2 - 1)K^{2n}\tilde{A}(n)^3 \\ &\quad + \{c_4(K^4 - 1) - 3c_2^2(K^2 - 1)\}K^{4n}\tilde{A}(n)^5 \\ &\quad - \{c_6(K^6 - 1) - 5c_2c_4(K^4 - 1) \\ &\quad + 3c_2(K^2 - 1)(4c_2^2 - c_4)\}K^{6n}\tilde{A}(n)^7 + \dots,\end{aligned}\quad (9)$$

which is a non-autonomous difference equation. In terms of the original variable x_n (see (6)), the renormalizaion equation (9) is transformed into an autonomous one.

$$\begin{aligned}x_{n+1} &= Kx_n - Kc_2(K^2 - 1)x_n^3 \\ &\quad + K\{c_4(K^4 - 1) - 3c_2^2(K^2 - 1)\}x_n^5 \\ &\quad - K\{c_6(K^6 - 1) - 5c_2c_4(K^4 - 1) \\ &\quad + 3c_2(K^2 - 1)(4c_2^2 - c_4)\}x_n^7 + \dots \\ &\equiv g_1(x_n),\end{aligned}\quad (10)$$

which gives the first branch g_1 of a return map on the unstable manifold: $x_{n+1} = g_j(x_n)$, where g_j is a multi-valued function of x_n and $j(= 1, 2, \dots)$ denotes a branch number. When $K \gg 1$ (strong chaos), it is easy to see that

$$g_1(x) = K[f(x) + \mathcal{O}(K^{-1})], \quad (11)$$

in general. If $(K - 1) \ll 1$ (weak chaos), we have,

$$g_1(x) = x + (K - 1)x(1 - x^2)^{1/2} + \mathcal{O}((K - 1)^2). \quad (12)$$

for the double well map for $K = 2.1$, an asymptotic expansion of g_1 truncated at x_n^7 is depicted in Fig .1. Although the truncated expression agrees well

with the exact numerical result for $x_n \lesssim 0.75$, it deviates considerably from the exact result near a singular point or a branch point ($x_n \approx 1.1$). In order to avoid this considerable discrepancy near a branch point, we restrict the domain of g_1 so that the return map g_1 is reversible, that is, $x_n \lesssim 0.75$. The reversible branch thus obtained is denoted by $\tilde{g}_1$. Then, from (2), we have the following functional map: $\tilde{g}_j \rightarrow \tilde{g}_{j+1}$

$$\tilde{g}_{j+1}(x_n) = 2x_n - \tilde{g}_j^{-1}(x_n) + \delta^2 f(x_n), \quad (13)$$

where the domain of x_n is chosen for each j so that a new branch $\tilde{g}_{j+1}$ is reversible. Using the functional map (13), we can construct each reversible branch of the return map $\tilde{g}_j$ from the first branch $\tilde{g}_1$ or g_1 step by step. As shown in the figures 2 and 3, the result agrees well with the exact one even if a truncated expansion of g_1 up to x_n^7 is used for $\tilde{g}_1$. Once the return map $\tilde{g}_j$ on the unstable manifold is constructed, the unstable manifold (x^u, p^u) is written as

$$p^u = x^u - \tilde{g}_j^{-1}(x^u), \quad (14)$$

because $p_n^u = x_n^u - x_{n-1}^u = x_n^u - \tilde{g}_j^{-1}(x^u)$.

Using the symmetry of the map (1):

$$x' = x - p, \quad p' = -p, \quad (15)$$

the stable manifold (x^s, p^s) is constructed as follows. The symmetry (15) gives

$$x^u = x^s - p^s, \quad p^u = -p^s. \quad (16)$$

From (16) and (14), we obtain the stable manifold in terms of $\tilde{g}_j$

$$p^s = x^s - \tilde{g}_j(x^s). \quad (17)$$

Therefore, the x-coordinate of a homoclinic point (x^h, p^h) is given by

$$x^h = \tilde{g}_i \tilde{g}_j(x^h). \quad (18)$$

Since the first intersection point (x^*, p^*) is located at $p^* = 0$ [4] on the first branch g_1 or $\tilde{g}_2$ in the present case, x^* is a fixed point of the first branch of the return map.

$$x^* = g_1(x^*) = \tilde{g}_2(x^*). \quad (19)$$

For $K \gg 1$ and $(K-1) \ll 1$, (11), (12) and (19) yield $x^* \approx \pm 1/\sqrt{2}$ and $x^* \approx \pm 1$ respectively.

3 Hénon map

Here, we construct an unstable manifold in the strange attractor of a dissipative dynamical system known as the Hénon map:

$$x_{n+1} = 1 - ax_n^2 + y_n, \quad (20)$$

$$y_{n+1} = bx_n, \quad (21)$$

where $a = 1.4$ and $b = 0.3$. This map has the hyperbolic fixed points:

$$\begin{aligned} x^* &= \frac{b - 1 \pm \{(b - 1)^2 + 4a\}^{1/2}}{2a}, \\ y^* &= bx^*. \end{aligned}$$

Let us consider the unstable manifold of one of the fixed points: $x^* = [b - 1 + \{(b - 1)^2 + 4a\}^{1/2}]/2a$, $y^* = bx^*$. This fixed point is known to be included in the closure of the strange attractor of the Hénon map. Setting $\hat{x}_n = x_n - x^*$ and eliminating the y -variable, we obtain the second-order difference equation from (20) and (21).

$$\hat{x}_{n+1} + 2ax^*\hat{x}_n - b\hat{x}_{n-1} = -a\hat{x}_n^2. \quad (22)$$

Near the origin of $\hat{x}$, we obtain the following formal series solution which vanishes as $n \rightarrow -\infty$.

$$\hat{x}_n = K^n[A - c_2A^2K^n + c_3A^3K^{2n} - c_4A^4K^{3n} + \dots],$$

where A is an arbitrary integral constant; $K = -ax^* + \{(ax^*)^2 + b\}^{1/2} \approx -1.923739$ is one of the eigenvalues of the linearized map of (22) and

$$\begin{aligned} c_2 &= \frac{aK^2}{D(K^2)}, & c_3 &= \frac{2a^2K^5}{D(K^3)D(K^2)}, \\ c_4 &= \frac{a^3K^8}{D(K^4)}\left\{\frac{1}{D(K^2)^2} + \frac{4K}{D(K^3)D(K^2)}\right\}, \\ D(K^j) &= K^{2j} + 2ax^*K^j - b. \end{aligned}$$

Following the same procedure as in the previous section, we renormalize A as

$$\begin{aligned} \tilde{A}(n) &= A - c_2A^2K^n + c_3A^3K^{2n} - c_4A^4K^{3n} + \dots, \\ \hat{x}_n &= \tilde{A}(n)K^n, \end{aligned} \quad (23)$$

and obtain a similar non-autonomous renormalization equation of $\tilde{A}$ as (9). Replacing $\tilde{A}(n)$ by $\hat{x}_n$ in the renormalization equation of $\tilde{A}$, we have

$$\begin{aligned}
\hat{x}_{n+1} &= K\hat{x}_n - c_2\hat{x}_n^2K(K-1) \\
&\quad + \hat{x}_n^3K(K-1)[c_3(K+1) - 2c_2^2] \\
&\quad + \hat{x}_n^4K(K-1)[-c_4(K^2 + K + 1) + 3c_3c_2(K+1) \\
&\quad \quad - c_2(5c_2^2 - c_3^3)] \\
&\quad + \hat{x}_n^5K(K-1)[c_5(K^3 + K^2 + K + 1) - 4c_2c_4(K^2 + K + 1) \\
&\quad \quad + 3c_3(3c_3^2 - c_3)(K+1) - 2c_2(7c_2^3 - 6c_2c_3 + c_4)] + \dots \\
&\equiv g_1(\hat{x}_n),
\end{aligned} \tag{24}$$

which gives the first branch g_1 of a return map on the unstable manifold: $\hat{x}_{n+1} = g_j(\hat{x}_n)$, where g_j is a multi-valued function of $\hat{x}_n$ and $j(= 1, 2, \dots)$ denotes a branch number. An asymptotic expansion of g_1 truncated at $\hat{x}_n^5$ is depicted in Fig. 4, where the truncated expression deviates considerably from the exact numerical result near $\hat{x}_n \approx 0.7$ and $\hat{x}_n \lesssim -1$. In order to avoid this considerable discrepancy, we restrict the domain of $g_1(\hat{x})$ to $-0.7 > \hat{x} > 0$, where the return map g_1 is reversible. The reversible branch in this way obtained is denoted by $\tilde{g}_1$. Then, from (22), we have the following functional map for neighboring reversible branches: $\tilde{g}_j \rightarrow \tilde{g}_{j+1}$

$$\tilde{g}_{j+1}(\hat{x}_n) = -a\hat{x}^2 - 2ax^*\hat{x}_n + b\tilde{g}_j^{-1}(\hat{x}_n), \tag{25}$$

where the domain of $\hat{x}_n$ is appropriately chosen for each j . Using the functional map (25), we construct each reversible branch of the return map $\tilde{g}_j$ from the first branch $\tilde{g}_1$ (24) step by step. Even when the asymptotic expansion of $\tilde{g}_1$ is truncated at $\hat{x}_n^5$, the result agrees well with the exact one as shown in Fig. 5 and Fig. 6. In Fig. 7, we see how well each branch of the return map constructed from an initial truncated map $\tilde{g}_1$ recovers the well-known many-leaved (fractal) structure of the strange attractor near the fixed point [9].

4 Conclusion

We construct asymptotic expansions for unstable manifolds of hyperbolic fixed points of the double-well map and the dissipative Hénon map, both of which exhibit the homoclinic chaos. Since the dimension of the manifold is

one, the dynamics on the manifolds is described by a return map with an infinite number of branches. An asymptotic expansion of the first branch of the return map is obtained by means of an updated renormalization method which is an extension of the RG method. We explicitly give a functional map, through which the other branches are obtained consecutively from the first branch. Thus a global asymptotic form of the unstable manifold is constructed to give the homoclinic chaos in the double-well map and the strange attractor in the dissipative Hénon map. Using a truncated expansion of the first branch of the return map, we calculate an approximate unstable manifold, which agrees well with the exact numerical result. A fixed point of the first branch of the return map is found to be the first homoclinic point in the double well map, and the approximate unstable manifold reproduces the well-known many-leaved (fractal) structure of the strange attractor in the Hénon map.

5 Acknowledgements

The authors would like to thank Y. Masutomi and other members of R-lab at Nagoya university for fruitful discussions.

References

- [1] H. Poincaré, Acta math. 13 (1890). 1
- [2] V.F. Lazutkin, I. G. Schachmannski and M. B. Tabanov, Physica 40 D (1989) 235.
- [3] K. Nakamura and M. Hamada J. Phys. A29 (1996) 7315.
- [4] V. G. Gelfreich, V. F. Lazutkin and N. V. Svanidze, Physica 71 D (1994) 82.
- [5] Y. Hirata, K. Nozaki and T. Konishi, Prog. Theo. Phys. 102-3 (1999) 701.
- [6] L. Y. Chen, N. Goldenfeld and Y. Oono, Phys. Rev. E 54 (1996) 376.
- [7] T. Kunihiro, J. Matsukidaira, Phys. Rev. E 57 (1998) 4817.

- [8] S. -i. Goto, Y. Masutomi, and K. Nozaki, Prog. Theo. Phys. 102-3 (1999) 471.
- [9] For example, H. G. Schuster, *Deterministic Chaos (An Introduction)*, Physik-Verlag, 1984.




Figure 1: The first branch of the return map on an unstable manifold for the double well map. While dots ($\cdot$) denote an exact solution calculated by means of numerical iterations, pluses ($+$) denote $\tilde{g}_1$ obtained by the renormalization method.




Figure 2: The exact return map on an unstable manifold for the double well map.




Figure 3: The return map plotted by means of matching the set of $\tilde{g}_j$.




Figure 4: The first branch of the return map on an unstable manifold for the double well map. While dots (·) denote an exact solution calculated by means of numerical iterations, pluses (+) denote $\tilde{g}_1$ obtained by the renormalization method.




Figure 5: The exact return map of the Hénon map's unstable manifold is plotted by numerical iterations.




Figure 6: The return map plotted by means of the matching the some of the $\tilde{g}_j$.




Figure 7: The return map of the Hénon map's unstable manifold near the hyperbolic fixed point $(\hat{x}_n, \hat{x}_{n+1}) = (0, 0)$ where the fractal structure appears. Pluses (+) express the exact map and line express the some of the $\tilde{g}_j$.